Contents

Community Evidence Summary iv
Executive summary v
Methodology vii

1.0 Introduction 1
1.1 Overview of Sight Loss 1
1.2 Causes of Sight Loss 2
1.3 Measure of Severity 3
1.4 The Detection of Sight Loss 4
1.5 Referrals, Diagnosis and Treatment 4
1.6 Preventable Eyesight Problems 5
1.7 The Economic Cost of Sight Loss 6
1.8 The Social Impact of Sight Loss 6
1.9 The International Context 7
1.10 The National and Local Context 9
1.11 Sight Loss in Birmingham 10
1.12 Registration and Certi cates of Vision Impairment (CVI) 10

2.0 Community Pro le 13

2.1 Getting the Best Start in Life 13
2.1.1 Prevalence of Sight Loss in children 13
2.1.2 Levels of Sight Loss in Children 13
2.1.3 Screening for Children 14
2.1.4 Growing up with Sight Loss 15

2.2 Mental Wellness and Balance 16
2.2.1 Mental Health 17
2.2.2 The Trauma of the Onset of Sight Loss 17
2.2.3 Dementia 18
2.2.4 The Emotional Cost of Living with Sight Loss 18
2.2.5 Provision of Support 19
2.2.6 Alcohol and Substance Abuse 20
2.2.7 Smoking 20
2.2.8 Alcohol and Substance Abuse 21
2.2.9 Domestic Abuse/Violence 21

2.3 Healthy and Affordable Food 22
2.3.1 Diet 22
2.3.2 Services to Support Healthy Lifestyles 22

2.4 Active at Every Age and Ability 23
2.4.1 Physical Exercise 23
2.4.2 Mobility and Transport 24
2.4.3 Sensory Coordination and Falls 25

2.5 Working and Learning Well 26
2.5.1 Education 27
2.5.2 Employment and Financial Well-being 27
2.5.3 Housing 29
2.5.4 Long-standing health impairment, illness or disability 29

2.6 Protect and Detect 30
2.6.1 Eyesight Screening 31
2.6.2 The Impact of COVID-19 on People with Sight Loss 33

2.7 Ageing Well and Dying Well 34
2.7.1 Diabetes 36
2.7.2 Strokes 36
2.7.3 Learning Disabilities 36
2.7.4 Hearing Impairment 36
2.7.6 Falls 36

2.8 Closing the Gaps 37
2.8.1 Life Expectancy 37
2.8.2 The Unequal Opportunities of People with Severe Sight Loss 38

2.9 Mitigating the Legacy of Covid-19 39

3.0 Conclusions 40

4.0 Appendices 41
Appendix 1: Availability of ECLO support by NHS Trust in Birmingham 41
Appendix 2: Unaddressed Visual Impairments 41
Appendix 3: Age distribution for Blind and Partially Sighted People in England 41
Appendix 4: Distribution of Certi cate of visual impairments across the UK 42
Appendix 5: Preventable Sight Loss Indicators 2019/20 42
Appendix 6: The Employment Status of People with Visual Impairment 42

5.0 Acknowledgements 43

6.0 References 44

List of Tables

Table 1:The eye diseases that cause severe visual impairment in the UK 2
Table 2A and 2B: The severity of visual impairment 3
Table 3:Current and Estimated Sight Loss Numbers in England and Birmingham 9
Table 4: Predicted Numbers Living with Sight Loss Conditions in Birmingham 12
Table 5: Estimated number of blind and partially sighted children by age group in Birmingham 14
Table 6: Sight Loss and Blindness in Older People by Eye Disease in England (2013) 35

List of gures

Figure 1: Unaddressed Visual Impairments 7
Figure 2: Age distribution for Blind and Partially Sighted People in England 10
Figure 3: Distribution of Certi cate of Visual Impairments Across the UK 11
Figure 4: Preventable Sight Loss Indicators 2019/20 11
Figure 5: The Employment Status of People with Visual Impairment 28

Executive Summary

Sight loss is a signi cant health issue that affects many people. In England, there were estimated to be 656,000 people with moderate to severe sight loss in 2021. In Birmingham, the estimate was 28,100. Sight loss can affect anyone at any stage of their life, and it is estimated that one in ve will start to live with sight loss during their lives. 50% of sight loss could have been treated if detected early. Screening for eyesight problems is recommended for everyone every two years, and major screening programmes are in place. In 2016/17, 12,995,512 NHS eyesight tests were conducted: 697,504 in Birmingham and the Black Country. This represents approximately 60% of the number of tests needed each year and leaves many people with possible eye defects that may deteriorate and become untreatable. In Birmingham, it is estimated that the total cost of sight loss, including direct and indirect costs, is £380,200 each year. Amongst the groups with the lowest take-up rates for screening are people with learning disabilities and autism, older people with dementia, people from deprived communities and people from Black and Minority Ethnic communities.

The major eye diseases that cause severe sight loss in the UK are AMD (Age-related Macular Degeneration), Cataracts, Glaucoma and Diabetic Retinopathy. There are 8,620 people registered blind or partially sighted in Birmingham, and 471 Certi cates of Visual Impairment were issued in 2016/17. People who are blind or partially sighted have great dif culty engaging in everyday community life and tend to lead isolated and sedentary lives: 40% say they are moderately or completely cut off from their local community. They tend to have lower attainment levels at school, with only one in four of working age being in employment. They also have major mobility problems both on foot and using public transport, and they often suffer nancial hardship. They have dif culty taking physical exercise and often have a poor diet, resulting in poor physical tness, which can lead to further eye problems. COVID has made isolation and loneliness an even greater problem for people with severe sight loss, and the incidence of mental health problems, especially depression, is high. 70% of people with severe sight loss are over 65, and the incidence of sight loss increases with age until it is 1 in 2 for the over 90s. Sight loss is frequently associated with other serious health problems, including learning disabilities and autism, dementia, diabetes and strokes, and people with severe sight loss have shorter life expectancy because of the link with other conditions.

The gaps and inequalities that this review reveals are:

- The gap in the number of eye-sight screening tests that are needed and are undertaken, with some communities, in particular, having low take-up rates.
- Problems of mobility on foot and on public transport mean that people with sight loss have dif culty engaging in community life.
- There are low levels of physical activity amongst people with sight loss and they have dif culty following a good diet. Together these factors lead to poor physical tness levels.
- People with sight loss have fewer opportunities to become self-suf cient and independent through educational attainment, employment, and nancial security.
- People with sight loss have low levels of emotional well-being and whilst there is support with medical care and physical rehabilitation, support for emotional wellbeing may be less available.

- Sight loss is associated with autism and dementia, and support for multiple long-term conditions may be fragmented.
- The elderly are most at risk of severe sight loss, resulting in them becoming more isolated, lonely, and more prone to mental health problems.
- People with sight loss in some communities are more at risk than others. These include: the most deprived areas, Black and Minority Ethnic communities, people with learning disabilities and autism and the elderly with dementia.
- The COVID pandemic has caused even greater isolation and loneliness and made access to health and community services even more dif cult for people with sight loss with consequential increases in mental health problems.



Methodology

Sight loss is one of the most prevalent health problems in the world. As a result, there is an extensive database on sight loss covering prevalence statistics and causes and treatments for every kind of sight loss impairment. To identify relevant sources, search terms were narrowly de ned and made speci c to each section of the report. Although some use was made of general databases, the main focus was databases specialized in sight loss. Apart from those included in the Introduction, all references to studies conducted outside the UK were excluded.

For the demographic section, the search terms used were ‘sight loss’ or ‘visual impairment’ and ‘statistics’ or ‘prevalence’ and ‘world’ or ‘global’ or ‘UK’ and ‘Birmingham’. Searches were undertaken using PubMed and Google Scholar and the specialist databases of:

- The World Health Organisation (https://www.who.int/)

1.3 Measure of Severity

Cataracts	A common eye condition prevalent in older people. The lens becomes less transparent and turns misty or cloudy. Cataracts, over time, can get worse and impact vision. A straightforward operation replaces the lens with an arti cial one. Numerous studies have demonstrated the cost bene ts of cataracts surgery in improving life quality and reducing the number of falls. In Birmingham, it's estimated that 8,640 people are living with cataracts. Between 2018 and 2030, there is estimated to be an increase of 19% in the number of people living with cataracts in Birmingham.
Glaucoma	A group of eye conditions in which the optic nerve can be damaged due to changes in eye pressure. Damage to sight can usually be minimised by early diagnosis and careful regular observation and treatment. Many glaucoma patients will attend regular appointments and take eye drops for the rest of their lives to prevent deterioration of vision. Some forms of glaucoma can be treated with laser surgery and surgery. RNIB estimates that 19,800 Birmingham citizens are living with ocular hypertension. A further 8,950 people are living with glaucoma. Between 2018 and 2030, there is estimated to be an increase of 16% in the number of people living with glaucoma in Birmingham.
Diabetic Eye Disease	People with diabetes are at risk of diabetic eye disease, which can affect the blood vessels in the eye. This can lead to permanent sight loss. Screening and early diagnosis with appropriate intervention are essential. RNIB estimates in Birmingham: • È£]Îää >`Õ•ÌÃ ...>Ûi Lii˜ `ˆ>}˜œÃi` ÜˆÌ... `ˆ>LiÌiÃ° • Ó£]Èää «iœ«•i >Ài •ˆÛˆ˜} ÜˆÌ... `ˆ>LiÌˆV ÀiÌ˜œ«>Ì...Þ° Of these, 1,990 have severe diabetic retinopathy, a later stage of the disease that is likely to result in signi cant and potentially certi able sight loss38. Between 2018 and 2030, there is estimated to be an increase of 8% in the number of people living with diabetic retinopathy in Birmingham

Source: WHO[1] and RNIB Sight Loss Data Tool v 4.3.1[2]

1.4 The Detection of Sight Loss

The World Health Organisation [5] estimates that globally at least 2.2 billion people have a near or distant vision impairment. In at least 1 billion (50%) of these cases, vision impairment could have been prevented or has yet to be addressed. The leading causes of vision impairment and blindness are uncorrected refractive errors and cataracts.

Prevention of avoidable sight loss is recognised as a key priority for the WHO's global initiative to eliminate avoidable blindness by 2020 – Vision 2020 – The Right to Sight [6], to which the UK is a signatory. This is also a key priority for Vision UK and is a particularly important issue in the ageing population.

The majority of people with vision impairment and blindness are over the age of 50 years; however, vision loss can affect people of all ages. It is estimated that, in 2020, 33.6 million people over the age of 50 were or could be classi ed as blind. 15 million (43%) of these cases were caused by cataracts [7]. In 2020, there were estimated to be 206 million people over 50 with mild or severe vision impairment globally. The leading cause is an uncorrected refractive error, with an estimated 86 million cases globally (42% of all cases).

Detecting sight loss must be undertaken at all stages of life and is a major industry in the developed world. There are 13 million NHS eye tests each year in England, leading to eight million outpatient appointments and 800,000 in patient procedures to treat and prevent sight loss [8].

The main causes of sight loss in the UK are:

- Uncorrected refracted error (39%)
- Age-related Macular Degeneration (AMD) (23%)
- Cataract (19%)
- Glaucoma (7%)
- Diabetic eye disease (5%)

1.5 Referrals, Diagnosis and Treatment

There are over eight million NHS outpatient appointments in eye clinics and ophthalmology departments in the UK every year and over 880,000 inpatient procedures (for cataracts and other surgical procedures) each year [9]. In Birmingham, in 2019/20, there were 397,835 outpatient appointments and 41,965 inpatient procedures [10]. The study [11] of ve communities that explored issues of accessing sight tests also explored the experiences of people being referred for treatment of glaucoma and diabetic retinopathy. The main problems identi ed were:

- Two-thirds of the people with sight loss are women.
- Adults with learning disabilities are ten times more likely to be blind or partially sighted.
- People on low incomes who, because of worries about costs, for example, the costs of spectacles, do not come forward for eyesight screening.
- People with diabetes are at risk of developing diabetic retinopathy.
- There are more than 25,000 blind and partially sighted children under 16 in the UK. It is estimated that half of these children have other disabilities[14, 15].

People from some ethnic communities are at greater risk of some of the leading causes of sight loss[16, 17]:

- Black African and Caribbean people are four to eight times more at risk of developing glaucoma compared to white people
- Within the Asian community in the UK, there is a much higher incidence of age-related cataracts when compared to white people
- The risk of diabetic eye disease is around three times greater in South Asian people compared to white people
- Black African and Caribbean people are also at a higher risk of diabetic171717

A BOLDER HEALTHIER BIRMINGHAM

There are considerable differences in the prevalence of sight impairment diseases across the globe. The prevalence of distance vision impairment in low- and middle-income regions is estimated to be four times higher than in high-income regions [33]. For near vision, unaddressed near vision impairment rates are estimated to be greater than 80% in Western, Eastern and Central Sub-Saharan Africa. In contrast, comparative rates in high-income regions of North America, Australasia, Western Europe, and Asia-Paci c are lower than 10%[34]. The prevalence of diseases across countries varies according to the availability of eye care services, their affordability, and the eye care literacy of the population. For example, the proportion of vision impairment attributable to cataracts is higher in low- and middle-income countries than in high-income countries. In high-income countries, diseases such as glaucoma and age-related macular degeneration (AMD) are more common. Among children, the causes of vision impairment vary considerably across countries. For example, congenital cataract is a leading cause in low-income countries, whereas, in middle-income countries, it is more likely to be retinopathy of prematurity. As in adult populations, the uncorrected refractive error remains a leading cause of vision impairment in all countries amongst children [35] [36].

The incidence and types of sight loss in the United Kingdom do not follow the dominant world pattern found in low- and middle-income countries. This report will focus on statistics and research on the forms of sight loss prevalent in the UK.

A systematic review[37] of progress in combating sight loss worldwide concluded that improvements in eye care services are succeeding in reducing the prevalence of profound sight loss. Still, rising populations and ageing populations will mean that the prevalence of severe eye disease is predicted to increase in the next 30 years. In particular, increases in the prevalence of two conditions associated with ageing, glaucoma and AMD (age-related macular degeneration), are predicted to increase signi cantly.

The RNIB forecast that there will be an increase of 17% in the number of people with severe sight loss in the UK by 2030 [38]. In Birmingham, the RNIB forecast that by 2030 there will be an increase of 18% in AMD cases, 19% in cataracts, 16% in glaucoma and 8% more cases of Diabetic Eye Disease[39].



Figure 3: Distribution of Certi cate of Visual Impairments Across the UK

Predicted eye disease and trends

Many people live with conditions that could lead to sight loss, and the number with these conditions is estimated to increase signi cantly in the next decade[66].

The estimated number of people who will be living with sight-threatening eye conditions in Birmingham from 2021 to 2030 [67] are shown in table 4.

Table 4: Predicted Numbers Living with Sight Loss Conditions in Birmingham

A statement of special educational needs (SEN) is issued to children to set out any additional help required in the education setting. In England in 2021, there were 16,555 pupils with a statement of special educational needs (SEN) or education, health and care (EHC) plans with vision impairment as their primary support need. Of these pupils, 8,486 were in primary school, 6,861 were in secondary school, and 1,208 were in special schools[71].

In Birmingham in 2021, there were 495 pupils with a statement of special educational needs (SEN) or education, health and care (EHC) plans with vision impairment as their primary support need. Of these pupils, 196 were in primary school, 200 were in secondary school, and 99 were in special schools[72].

In Birmingham[73], in 2021, there were an estimated 550 blind and partially sighted children aged 0-16 and 350 blind and partially sighted young people aged 17-25.

Table 5: Estimated number of blind and partially sighted children by age group in Birmingham

Age group	Blind	Partially sighted	Total
0 to 16 years	140	410	550
17 to 25 years	90	260	350

study of screening in 44 special schools in Wales found that 42% of pupils had received no screening[80]. When the screening was undertaken, 51% had some abnormality that warranted action to prevent risk to sight.

In Birmingham, the school entry vision screening service undertakes an eye test for every child during their reception year [81]. It is offered in every primary school in Birmingham, including special schools and religious schools. At this test, the child's eyes are tested for squints, coordination between the two eyes, and problems with vision.

2.1.4 Growing up with Sight Loss

There is a higher prevalence of children with visual impairment being

• •••• • - • €- • ••
,ƒ„ , €ƒ € ••„„ , , …
†• †•• , ,ƒ-•‡ • •
• • , ƒ … •• „ ‡•ƒ …•

„ƒ † ,• ƒ„ƒ••

People with poor eyesight reported that discrimination had a negative effect on their wellbeing, contributing to loneliness, depression and lower levels of life satisfaction

2.2.5 Provision of Support

The RNIB survey of 1,200 people with sight loss found that only 17% reported being offered any emotional support [110]. Despite many different

2.2.8 Alcohol and Substance Abuse

Excessive alcohol consumption and illicit drugs can have a very damaging effect on ocular functions. A meta-analysis of studies of the relationship between alcohol consumption and the development of cataracts, for example, concluded that heavy drinking, de ned as 20g per day or more, was associated with cataract development but that there was no association with more moderate drinking [124].

There is very little evidence that people with severe visual impairments in the UK drink alcohol excessively or have an illicit drug habit. There is some evidence that in the process of adjusting to a major loss of visual functioning, some people may resort to alcohol or drugs as coping mechanisms. A qualitative study[125] of users' perspectives on substance use found that people with severe visual impairment thought that alcohol led to their sight loss or contributed to it. That substance use helped them to cope with the consequences of sight loss. A qualitative study of 30 male [126] and 9 female[127] service personnel who had sustained a vision impairment found that some personnel used alcohol and drugs as a coping mechanism when adequate support to adjust to a new lifestyle was absent.

2.2.9 Domestic Abuse/Violence

No statistics are available that detail how people with severe sight loss are subjected to abuse either in the home or when in public places. However, the RNIB[128] reports many cases where people with severe sight loss have been abused by partners upon whom they depend for care and others who have been assaulted or robbed whilst travelling. People with severe sight loss report feeling vulnerable and defenceless in many circumstances.

2.3 Healthy and Affordable Food

2.3.1 Diet

A poor diet has been identi ed as a risk factor in both the onset and subsequent management of several major visual impairment causes in the UK. A systematic review of the impact of visual impairment on nutritional status[129] concluded that people in the UK with severe visual impairments tended to be obese, i.e., to have an abnormal BMI index over 30. Still, there were also reported cases of malnutrition. A strong correlation has also been found between type II diabetes and poor diet, and people with diabetes are likely to develop diabetic retinopathy.

A poor diet is a risk factor concerning all major causes of visual impairment in the UK: age-related macular degeneration, diabetic retinopathy, cataract and glaucoma. A review of dietary factors [130] concluded that the risk might be decreased by a diet that increases the intake of green leafy vegetables and reduces the glycaemic index by, for example, increasing the intake of sh that contain Omega-3 fatty acids. Vitamin supplements can also reduce the risk.

People with severe visual impairments face many challenges that make it dif cult to sustain a healthy diet. Mobility problems may make shopping dif cult and preparing and cooking food may be additional burdens [131].

2.3.2 Services to Support Healthy Lifestyles

The Birmingham and Solihull Health and Sight Loss Evidence Base[132] lists the services that help people improve their diet, stop smoking and reduce dependence on alcohol and drug consumption. However, it is noted thatmyac <</Lang (en-GB)/MCID actors

%‰

2.4 Active at Every Age and Ability

Active at Every Age and Ability Key Findings:

- People with severe Sight Loss partake in much less physical

2.4.3 Sensory Coordination and Falls

Many people with severe sight loss do not have the con dence to venture out to engage in physical exercise because they fear they may be involved in an accident or may have a fall. A survey of the hospital records in the UK in 1999 found that of the 2.35 million recorded falls, 189,000 (8%) occurred to people with a visual impairment, of which 89,500 were attributed to the visual impairment itself[143].

The RNIB estimates that in Birmingham, 3,270 people with sight loss over 65 experience a fall per year and that, of these falls,1,550 are directly attributable to sight loss. 260 people over 65 with sight loss experience a severe fall per year that results in hospital admission through A&E. Of these severe falls,120 are directly attributable to sight loss [144].

People with little or no sight have lower coordination and balance levels than the sighted. Physical exercises involving perceptual motor skills can be dif cult and potentially dangerous [145]. It is one of the key reasons people with severe sight loss are likely to do so with less vigour than sighted people, even when taking physical exercise.

People with severe sight loss often have other serious health issues, and the combination of conditions may make taking physical exercise dif cult. The other conditions associated with severe sight loss include learning disabilities, autism, dementia, diabetes, and hearing loss [146].

Engaging in regular physical exercise in their own homes may be an option for people with severe sight loss. Still, many, especially in deprived areas, do not have the facilities for it, and when it is introduced, it may not be sustained [147].

One of the reasons people with severe sight loss give for not being able to take regular physical exercise is that they need help, particularly with walking and transport. Ideally, the help would be from a sighted person. However, in the RNIB My Voice survey of 1,200 people with severe visual impairments, 19% of respondents reported a lack of sighted help or assistance with transport [148].

People with severe sight loss express a strong desire to be as independent as possible. There are many aids and services designed to enable them to be independent in walking, travelling, and physical exercise. These include the traditional white canes, guide dogs and, more recently, technology aids that, for example, use GPS to identify the location of the person and download map information to braille readers [149]. Although there are no available statistics on the use of these aids by people with severe sight loss, it is probable that the percentage of people who access these services is quite low. People would need to know of these aids and services and then be able to locate them, and it can be dif cult for people with recent severe sight loss to access support services. The RNIB report that 49% of people in contact with their local authority do not go on to receive an assessment for vision rehabilitation support [150].

There are estimated to be 4,800 working guide dogs in England, serving an estimated population of 245,000 blind people, and there are long waiting lists[151].

Although digital technology aids are increasingly sophisticated and available, people with severe sight loss are less able to use them than sighted people. The RNIB My Voice survey[152] found that 82% of younger people aged 18-29 with sight loss were making wide-ranging use of digital technology compared with 25% of people over 75 years old. However, it is estimated that 96% of the younger age group who are sighted now make regular use of digital technology. In the survey, respondents reported that sight loss remained a barrier to using digital technology to support travel and physical exercise. Although there were many aids to help people with sight loss make use of the technology, they were more usable in the home and in employment than whilst travelling.

2.5.1 Education

The RNIB My Voice survey of 1,200 people with severe sight loss found that 27% had a degree or higher-level quali cation, while 33% reported no quali cations. Older participants were more likely to have no quali cations than younger people. 38% of people of pension age reported having no quali cations compared to 9% of people under 50 years [153]. Given that the onset of slight loss may be at any stage of life, most quali cations may be gained before people suffer sight loss. However, the question of how people who have sight loss cope in the education system will be signi cant for many people and is addressed below.

At both primary school and secondary school levels, pupils with visual impairments record lower levels of achievement than their sighted counterparts. However, 50% of pupils with visual impairment also have other special educational needs. An analysis for the RNIB[154] of the National Pupil Datasets for 2008/09 held by the Department of Education examined the performance of sighted pupils, pupils with only a visual impairment and pupils with a visual impairment and other special educational needs. At all stages of primary education, the sighted pupils did best, followed by those with only a visual impairment and those with another particular educational need. These gaps widened with every year of education. This pattern was repeated at the age of 16 when pupils undertook GCSE examinations

There are no comparable studies of attainment levels in higher education. However, there is evidence that the challenges of studying in universities and colleges mean that visually impaired and blind students represent a relatively small percentage of the total number of students. An analysis of Higher Education Statistics Agency data for 2017/2018 [155] found 3,170 students registered and identi ed as visually impaired across all levels of study. A number of studies [156] [157] have followed the progress of students with severe visual impairments and have identi ed the following challenges:

Getting support to study

Universities and colleges are required under the UK Equality Act 2010 to make reasonable adjustments for students with special needs, including those with sight loss. Many now have provisions including special computer equipment, help with notetaking, special arrangements for examinations, and extra time. Each student has unique needs, and most universities and colleges have a disablement of cer to help organise appropriate assistance. Despite this progress, most studies nd that assistance is patchy and that students struggle to maintain the progress made by sighted colleagues.

Living independently away from home

Most students move to live on or near a university campus to study, which means they have the challenge of living independently, usually for the rst time. The studies have found that students are often not well prepared for this new life. A common recommendation is that in transitioning from secondary school to higher education, in addition to their academic studies, pupils should have detailed training in living independently.

2.5.2 Employment and Financial Well-being

Estimates of the employment rate of people with severe sight loss who are of working age vary from 25%[158] to 33%[159]. All studies agree that the employment rate is signi cantly below the UK rate for the general population of 4% (8% in Birmingham)[160]. A study of 1,000 people in England with severe sight loss aged between 18 and 64 provided the following statistics on employment rates and other outcomes as shown by gure 5 (see Appendix 6 for full data) [161].

2.6.1 Eyesight Screening

Screening for eyesight problems is essential if corrective measures are put in place. Eyesight problems can occur at any stage of life and are particularly prevalent in older age groups. Regular eye checks are necessary throughout life, and the recommendation is that eye checks are made at least every two years[172].

The RNIB estimates that 50% of blindness and severe sight loss cases could be prevented [173] [174] if detected and treated in time. This is mainly due to uncorrected refractive error and untreated cataracts. Low take-up of sight tests can lead to detection after the condition has deteriorated, and intervention may be less effective.

Prevention of avoidable sight loss is recognised as a key priority for the WHO's global initiative to eliminate avoidable blindness by 2020 – Vision 2020 – The Right to Sight[175], to which the UK is a signatory and is also a key priority for Vision UK.

The Availability and Take-Up of Eyesight Tests

Free NHS sight tests are available to people in England who meet certain criteria. For example, people aged 60 or over, children aged 15 and under, and people receiving certain bene ts. Public data on sight tests only records those paid for by the NHS. Data about privately funded sight tests are not available.

The RNIB Top((1rtBir)12795,512NHS eyesight tests were conducted in England in 2016/17, of which 697,504 were in Birmingham and the Black Country area[176] [177]. In Birmingham, in terms of the groups eligible for NHS sight tests:

/Span<</ActualText<FEFF0020>>BDC /P9.685 129.0699 Tm (sight tests:)Tj ET EM759 Tm (•)Tj

•

- People with autism and learning disabilities
- Older people, particularly those with dementia and with mobility problems
- Working-age males
- People from areas of multiple deprivations
- Black and ethnic minority communities

A study[183] that explored the reasons for not attending regular eyesight tests in ve UK communities with large ethnic communities and/or high levels of deprivation focused on exploring issues concerning speci c eye conditions in each community. The study included 34 focus groups (289 participants), 56 case study interviews with members of the target populations and 55 interviews with eyesight test providers

The ve local communities were:

- Bradford (diabetic retinopathy in a Pakistani population experiencing deprivation, aged 40 to 65)
- Cwm Taf (glaucoma in a white, deprived population, aged 40 years and above)
- Glasgow (diabetic retinopathy in Pakistani people living in af uent and deprived areas aged 40 to 65)
- Hackney (glaucoma in a Caribbean population, aged 40 to 65)
- West Belfast (glaucoma in a white, disadvantaged population, aged 40 to 65).

The reasons given in all communities for not attending regular eyesight tests included:

1. Limited community awareness of eye health - Eyesight was not regarded as a health issue requiring regular assessment.
2. Symptom-led demand for eye examinations - People only went for an eyesight test when they became aware that their eyesight was deteriorating.
3. The cost and retail element associated with primary care - Most opticians were located in the high street, often within large retail chains, and people associated them with selling expensive spectacles. They were more inclined to view independent opticians as professional health specialists, whereas opticians in large chains were regarded more as sales agents.
4. Community-speci c reasons - The language was a barrier to accessing eye health services for some people in Glasgow. Physical access was an issue in Cwm Taf, re ecting the particular geography and relatively poor transport links of the Welsh Valleys.

Consultations on Eye Health

In 2013, visual health was made a clinical priority for GP practices. The England Vis Sn9.898tions 184 <</Lang (en-GB)/MCID 2163 >>BDC BT /T 0 0 11 465.

The audit identi ed ve barriers:

- Physical barriers – mobility issues making travel to the practice dif cult
- Communication barriers – inability to read and understand practice notices and calling systems
- Procedural barriers – dif culties with paperwork and computerised check-in systems
- Medical ob.026f-e1.0236 403.7471 Tm [(Med)-1 (ical ob.0260f h0a(Med)-1 (icqga0ang (8 (11 51.0236 423.4164 Tm (in systems)T03.7471 Tm (iputesigns of iP <<



2.7 Ageing Well and Dying Well

Ageing and Dying Well Key Findings:

- In the UK in 2019/20 69.9% of people with severe Sight Loss were aged 65 or over.
- In Birmingham in 2019/20 69.8% of people with severe sight loss were aged 65 or over.
- Sight Loss increases with age. 1 in 8 people over 60 have Sight Loss, this rises to 1 in 2 in those aged over 90.
- The leading causes of sight loss in older people are refractive errors, AMD (age-related macular degeneration), cataracts, glaucoma and diabetic retinopathy.
- There are nearly double the number of females over 65 with Sight Loss compared with males.
- Older people with Sight Loss frequently also have other major health conditions such as hearing loss, dementia, strokes, learning disabilities, disbetes, physical disabilities and poor mental health.
- Older people with Sight Loss have a high incidence of falls leading to hospitalisation.
- There are no agreed statistics on the life expectancy of older people with sight loss but hazard ratios range from 1:1.29 for mild Sight Loss to 1:89 for severe Sight Loss.
- The reduced life expectancy of people with Sight Loss is mostly attributable to the life expectancy of associated health conditions.

2.7.1 Diabetes

Half of all people with diabetes in the United Kingdom are over 65 years



2.8 Closing the Gaps

2.8.1 Life Expectancy

Although there are no statistics for the life expectancy of people with sight loss in the UK, several studies have established that life expectancy for people with sight loss is signi cantly shorter than for people who have no sight loss. In a systematic review of 17 studies of sight loss and mortality,[198] evidence showed that life expectancy decreased as the severity of the sight loss increased. Expressed as a Hazard Ratio, the greatest mortality risk was amongst those with severe vision impairment or blindness (1:1.89), compared with less risk of mortality for those with mild impairment (1:1.29).

A review of 13,569 patients from 49 UK GP practices[199] examined the hazard ratios for the different eye diseases. It found that for AMD, there was a hazard ratio of 1.40, and for cataracts, 1.35. However, after removing the

Amongst the associations with other conditions, the lower life expectancy of people with learning disabilities has been noted in section 8. Diabetes UK estimate[202] that people with Type 1 diabetes may have their life expectancy reduced by 20 years, and for people with Type II diabetes, the reduction may be up to 10 years. Life expectancy for people who have had strokes may be shortened by 5 to 10 years depending on the type of stroke: 1 in 4 people die within one year of having a stroke [203].

2.8.2 The Unequal Opportunities of People with Severe Sight Loss

Sight loss is a broad category that extends from people who have a temporary visual impairment to those living with a permanent incapacity that seriously inhibits their life opportunities. Most people have a temporary sight problem; for example, they need a refractive error-correcting, and if they are prescribed suitable spectacles, they have 'normal' sight once again. However, a signi cant group at the other end of the spectrum has severe sight loss, are blind or have partial sight, and has to live with these disabling conditions. On a wide variety of fronts, the second group have much less opportunity in life than sighted people. There are considerable gaps in what people with severe sight loss can achieve in society compared with people with normal sight. These inequalities can be found in the areas discussed below:

- Whilst educational opportunities at primary school for blind or partially sighted children enable them to make a good start in education, these children steadily drop behind in attainment through secondary and tertiary education. There are relatively few blind and partially sighted students at the university and college level that achieve a higher education award.
- People who are blind or partially sighted tend to become isolated and lonely in their homes and experience many mobility problems when leaving home. Few of them are in employment, and they may experience severe nancial dif culties. These factors contribute to a high level of mental health problems.
- To engage in the world as sighted people do, blind or partially sighted people need support. They may receive appropriate medical support to manage their condition, but the social care structure that supports their wider life needs is fragmented and slow. Too often, there is no coordinated support plan, and support relies on friends and families.
- People with severe and permanent sight loss usually have other serious health conditions, notably learning disorders, diabetes and dementia. Apart from the dif culties of managing several conditions, the other conditions also make it more dif cult for the issues of sight loss to be addressed; for example, a learning disability or dementia may make it more dif cult for a person who is blind or partially sighted to engage with the medical profession: to attend screening sessions, manage medication, achieve greater mobility etc.
- The majority of severe sight loss occurs in old age and becomes more prevalent with advanced old age. It can have a very harmful effect on the quality of life in old age, leading to even more isolation, loneliness and depression. In a complex way, the very common association with other serious health conditions, for example, strokes and dementia, can also lead to reduced life expectancy.

Although all the above inequalities can affect anybody with severe sight loss, the impacts are particularly extreme for the Black and Minority Ethnic communities and people in areas of multiple deprivations. People in these communities with severe sight loss have additional problems navigating the complex health and social care system to receive the help and support they need.

2.9 Mitigating the Legacy of Covid-19

The lockdowns and other restrictions during the pandemic have led to many people experiencing loneliness and poor mental health. People with severe sight loss already face these deprivations, and, although it is dif cult to quantify the impact, the pandemic has likely deepened their experience of isolation from the rest of the community.



3.0 Conclusions

The scale of sight loss and the fact that it can occur at any stage in life makes it a major public health challenge for any nation. As a developed nation, the UK has made major strides in treating many sight loss conditions and has an extensive health and social care system to detect, treat and manage the condition. Nevertheless, there remain many challenges to address, and for people with severe sight loss, life can be very restrictive, and wellbeing can be very poor. Seventy per cent of sight loss develops in the elderly. Because the UK has an increasingly ageing population, there is predicted to be a signi cantly increasing demand for sight loss in the health and social system.

Approximately 50% of sight loss goes undetected at an early stage and can lead to serious sight problems at a later stage. A major challenge is reaching the communities with low take up rates for regular screening.

For the people with severe sight loss who are blind or partially sighted, life can be very restricted, well-being can be hard to achieve, and life expectancy can be limited. Many become isolated and lonely, and mobility problems can restrict access to employment, education, physical activity and other kinds of engagement with normal community life. People with severe sight loss are also likely to have other serious health concerns, and the majority are elderly. A key objective needs to strengthen the current health and social system to provide a comprehensive and timely support system for this vulnerable population.



4.0 Appendix

Appendix 1: Availability of ECLO support by NHS Trust in Birmingham

Birmingham and the Black Country Area Team	Ophthalmic outpatient appointments 2019/20	RNIB-trained ECLOs (Jan 2021)
Birmingham Women's And Children's NHS Foundation Trust	10,745	No
Sandwell and West Birmingham Hospitals NHS Trust	142,310	Yes
The Dudley Group NHS Foundation Trust	35,875	Yes
The Royal Wolverhampton NHS Trust	74,695	Other forms of support
University Hospitals Birmingham NHS Foundation Trust	121,385	Yes
Walsall Healthcare NHS Trust	12,825	No

Source: RNIB Sight Loss Data Tool v4.3.1: Birmingham

Appendix 2: Unaddressed Visual Impairments

Visual Impairment Categories	Cases (Millions)
Uncorrected Presbyopia	826
Unaddressed Refractive Error	123.7
Cataract	65.2
Glaucoma	6.9
Corneal Opacities	4.2
Diabetic Retinopathy	3
Trachoma	2

Appendix 3: Age distribution for Blind and Partially Sighted People in England

Age	Partial sight/Sight impaired persons	Blind/Severely sight impaired persons
0 to 4	470	520
5 to 17	5,145	4,010
18 to 49	18,030	19,020
50 to 64	16,585	19,010
65 to 74	14,710	15,160
75+	85,445	78,175
Totals	140,390	136,300

Appendix 4: Distribution of Certi cate of visual impairments across the UK

Country	Distribution of CVIs
England	83%
Wales	5%
Scotland	10%
N. Ireland	2%

Appendix 5: Preventable Sight Loss Indicators 2019/20

PH Outcomes Framework Indicator	Birmingham (Proportion per 100,000)	England (Proportion per 100,000)
E12d Sight Loss Certi cation	37.4	41.4
E12c Diabetic Eye Disease	4.6	2.9
E12b Glaucoma 40+	9.2	12.9
E12a AMD 65+	98.4	105

Appendix 6: The Employment Status of People with Visual Impairment

Age	Employment	Unemployment	Student	Retired	Long Term Sick or Disabled
18-29	33%	22%	22%	0%	17%
30-49	44%	13%	1%	4%	32%
50-64	22%	9%	0%	19%	45%

5.0 Acknowledgements

Contributors:

Ricky Bhandal –Service Lead, Birmingham City Council

Caroline Chioto –Senior Of cer, Birmingham City Council

Alice Spearing – Senior Of cer, Birmingham City Council

Joseph Merriman – Senior Of cer, Birmingham City Council

Rhys Boyer – Of cer, Birmingham City Council

Alexander Robinson – Support Of cer, Birmingham City Council

Jordan Francis – Graduate Of cer, Birmingham City Council

Tariro Mandisodza - Graduate Of cer, Birmingham City Council

Manuela Engelbert - Graduate Of cer, Birmingham City Council

Nazmin Khanom - Graduate Of cer, Birmingham City Council

Dr. Justin Varney – Director of Public Health, Birmingham City Council

Tessa Lind eld – Assistant Director of Public Health, Birmingham City Council

Modupe Omonijo - Assistant Director of Public Health, Birmingham City Council

Alan Davies – Head of Marketing, Birmingham City Council

Carl Madden – Designer, Birmingham City Council

Barques Design – Production of Artwork/Typesetting

6.0 References

1 World Health Organisation (2019) WHO ‘World report on vision’. Geneva: World Health Organization; 2019. License: CC BY-NC-SA 3.0 IGO. SBN: 9789241516570

2 RNIB (2021) Sight Loss Data Tool Version 4.3.1: Birmingham 2021

3 Vision Loss Expert Group of the Global Burden of Disease Study (2020) Causes of blindness and vision impairment in 2020 and trends over 30 years: evaluating the prevalence of avoidable blindness in relation to “VISION 2020: the Right to Sight”. Lancet Global Health 2020. doi.org/10.1016/S2214-109X(20)30489-7

4 Vision Loss Expert Group of the Global Burden of Disease Study (2020) Causes of blindness and vision impairment in 2020 and trends over 30 years: evaluating the prevalence of avoidable blindness in relation to “VISION 2020: the Right to Sight”. Lancet Global Health 2020. doi.org/10.1016/S2214-109X(20)30489-7

5 World Health Organisation (2019) WHO ‘World report on vision’. Geneva: World Health Organization; 2019. License: CC BY-NC-SA 3.0 IGO. SBN: 9789241516570

6 Vision 2020 The International Agency for the Prevention of Blindness https://www.iapb.org/vision-2020/ Accessed 15/02/22

7 Vision Loss Expert Group of the Global Burden of Disease Study. (2020) Trends in prevalence of blindness and distance and near vision impairment over 30 years: an analysis for the Global Burden of Disease Study. Lancet Global Health 2020. doi: 10.1016/S2214-109X(20)30425-33.

8 RNIB (2021) Sight Loss Data Tool Version 4.3.1: England 2021

9 Slade J. and Edwards R (2015) MyVoice 2015: The views and experiences of blind and partially sighted people in the UK. RNIB.

10 Vision Loss Expert Group of the Global Burden of Disease Study. (2020) Trends in prevalence of blindness and distance and near vision impairment over 30 years: an analysis for the Global Burden of Disease Study. Lancet Global Health 2020. doi: 10.1016/S2214-109X(20)30425-33.

11 Fricke, TR, Tahhan N, Resnikoff S, Papas E, Burnett A, Suit MH, Naduvilath T, Naidoo K, (2018) Global Prevalence of Presbyopia and Vision Impairment from Uncorrected Presbyopia: Systematic Review, Meta-analysis, and Modelling, Ophthalmology. 2018 May 9

12 Scase MO and Johnson MR (2005) Visual Impairment in Ethnic Minorities in the UK International Congress Series 1282 438-442

13 Scase MO and Johnson MR (2005) Visual Impairment in Ethnic Minorities in the UK International Congress Series 1282 438-442

14

32 Fricke, TR, Tahhan N, Resnikoff S, Papas E, Burnett A, Suit MH, Naduvilath T, Naidoo K, (2018) Global Prevalence of Presbyopia and Vision Impairment from Uncorrected Presbyopia: Systematic Review, Meta-analysis, and Modelling, Ophthalmology. 2018 May 9

33 World Health Organisation (2019) WHO 'World report on vision'. Geneva: World Health Organization; 2019. License: CC BY-NC-SA 3.0 IGO. SBN: 9789241516570

34 Vision Loss Expert Group of the Global Burden of Disease Study (2020) Causes of blindness and vision impairment in 2020 and trends over 30 years: evaluating the prevalence of avoidable blindness in relation to "VISION 2020: the Right to Sight". Lancet Global Health 2020. doi. org/10.1016/S2214-109X(20)30489-7

35 Vision Loss Expert Group of the Global Burden of Disease Study (2020) Causes of blindness and vision impairment in 2020 and trends over 30 years: evaluating the prevalence of avoidable blindness in relation to "VISION 2020: the Right to Sight". Lancet Global Health 2020. doi. org/10.1016/S2214-109X(20)30489-7

36 Vision Loss Expert Group of the Global Burden of Disease Study. (2020) Trends in prevalence of blindness and distance and near vision impairment over 30 years: an analysis for the Global Burden of Disease Study. Lancet Global Health 2020. doi: 10.1016/S2214-109X(20)30425-33

37 Vision Loss Expert Group of the Global Burden of Disease Study (2020) Causes of blindness and vision impairment in 2020 and trends over 30 years: evaluating the prevalence of avoidable blindness in relation to "VISION 2020: the Right to Sight". Lancet Global Health 2020. doi. org/10.1016/S2214-109X(20)30489-7

38 RNIB (2018) Eye Health and Sight Loss Stats and Facts April 2018

39 RNIB (2021) Sight Loss Data Tool Version 4.3.1: Birmingham 2021

40 Pezzullo et al (2019) The economic impact of sight loss and blindness in the UK adult population. RNIB and Deloitte Access Economics. Prevalences applied to subnational population projections

41 International Agency for the Prevention of Blindness (2021) Country Map & Estimates of Vision Loss: United Kingdom https://www.iapb.org/learn/vision-atlas/magnitude-and-projections/countries/united-kingdom/

42 Galvani S., Livingston W. and Morgan Hobal B Tm (41)Tj ET EMC /P <</Lang (en-GB)/MCID 2923 >>BDC BT /Span<</ActualText<FEFF0009>>> BDC 8 0 0 8 48.581 226.015 Tm ( )Tj EMC 1.014 0 Td [(Inter)-18 (na (na

102 Gosney M, Victor CR and Nyman SR (2010) Emotional Support to People with Sight Loss Thomas Pocklington Trust Occasional Paper April 2010 No 26.

103 England Vision Strategy (2018) Birmingham and Solihull Eye Health and Sight Loss Evidence, England Vision Strategy, NHS England.

104 Cumberland P.M. & Rahi J. (2016) Visual function, social position, and health and life chances: The UK Biobank study JAMA Ophthalmol. 2016;134(9):959-966.

105 Lawrence V. and Murray J. (2010) Balancing independence and safety: the challenge of supporting older people with dementia and sight loss Age and Ageing 2010; 39: 476–480.

106 RNIB (2018) Eye health and sight loss stats and facts RNIB April 2018

107 Swanson A, Edwards R (2015) MyVoice 2015: The views and experiences of blind and partially sighted people in the UK. RNIB.

108 World Health Organization (2019) World report on vision. Geneva: World Health Organization; 2019. License: CC BY-NC-SA 3.0 IGO. SBN: 9789241516570

138 Aslan U. Calik B and Kiti A (2012) The effect of gender and level of vision on the physical activity level of children and adolescents with visual impairment Research in Developmental Disabilities, 1799-1804, 33(6).

139 Slade J and Edwards R (2015) MyVoice 2015: The views and experiences of blind and partially sighted people in the UK. RNIB.

140 Slade J and Edwards R (2015) MyVoice 2015: The views and experiences of blind and partially sighted people in the UK. RNIB.

141 Slade J and Edwards R (2015) MyVoice 2015: The views and experiences of blind and partially sighted people in the UK. RNIB. 261 (.iapb.or)18 (g/)]TJ ET E:31.3t<FETj ng (en-GB)/MCID 3241 >>BDC BT 8 0 0 8 56.6929 419.0229 Tm [(sighted people in 6)-0.9 sighted people in the UK. RNIB. 265 BT 8 0 0 8 56.6929 385.519 Tm (144)Tj ET EMC /P <</Lang (en-GB)/MCID 3249 >>267 /P04 Slade J and Edwards R (2015) MyVoice 2015: The views and experiences of blind and pa6.1 (.iapb.or)18 (g/)]TJ ET E:/

142 English Vision Strategy (2018) Birmingham and Solihull Eye Health and Sight Loss Evidence Base NHS 2018.

143 Scuffham P., Chaplin S. and Legood R. (2003) Incidence and costs of unintentional falls in older people in the United Kingdom J Epidemiol Community Health 2003;57:740–744.

144 RNIB (2021) RNIB Sight Loss Data Tool v 4.3 Birmingham

145 Vision 2020 The International Agency for the Prevention of Blindness https://www.iapb.org/vision-2020/

018-2836-0

175 Vision 2020 The International Agency for the Prevention of Blindness https://www.iapb.org/vision-2020/ Accessed 15/02/22

176 Vision 2020 The International Agency for the Prevention of Blindness https://www.iapb.org/vision-2020/ Accessed 15/02/22

177 Vision Loss Expert Group of the Global Burden of Disease Study. (2020) Trends in prevalence of blindness and distance and near vision impairment over 30 years: an analysis for the Global Burden of Disease Study. Lancet Global Health 2020. doi: 10.1016/S2214-109X(20)30425-33.

178 England Vision Strategy (2018) Birmingham and Solihull Eye Health and Sight Loss Evidence Base 2018

179 England Vision Strategy (2018) Birmingham and Solihull Eye Health and Sight Loss Evidence Base 2018

180 England Vision Strategy (2018) Birmingham and Solihull Eye Health and Sight Loss Evidence Base 2018

181 England Vision Strategy (2018) Birmingham and Solihull Eye Health and Sight Loss Evidence Base 2018.

182 RNIB (2018) Eye Health and Sight Loss Stats and Facts RNIB 2018

183 Hayden, C et al (2012) The barriers and enablers that affect access to primary and secondary eye care services across England, Wales, Scotland and Northern Ireland. RNIB report: RNIB/CEP/IR/01, 2012.

184 England Vision Strategy (2018) Birmingham and Solihull Eye Health and Sight Loss Evidence Base 2018

185 Focus Birmingham (2017) Better Healthcare Access for People with Visual Impairment https://www.vistablind.org.uk/documents/246/Better_Healthcare_Access_For_People_With_Visual_Impairment._nal_report....pdf

186 Ting, D.S.J., Krause, S., Said, D.G. et al. (2021) Psychosocial impact of COVID-19 pandemic lockdown on people living with eye diseases in the UK. Eye 35, 2064–2066 (2021). https://doi.org/10.1038/s41433-020-01130-4 .

187 RNIB (2021) Voice of the Customer: Living with Sight Loss through Lockdown in the United Kingdom (UK) RNIB 2021.

188 RNIB (2018) Eye Health and Sight Loss Key Stats and Facts RNIB 2018

189 Deloitte Access Economics (2014) The economic impact of sight loss and blindness in the UK adult